



FACULDADE DE MEDICINA
UNIVERSIDADE DO PORTO

MESTRADO INTEGRADO EM MEDICINA

2013/2014

Mónica Sofia Gonçalves Garrido

SOX2 and CDX2 in gastric intestinal metaplasia
and dysplasia: impact on differentiation

março, 2014

FMUP

Mónica Sofia Gonçalves Garrido

SOX2 and CDX2 in gastric intestinal metaplasia
and dysplasia: impact on differentiation

Mestrado Integrado em Medicina

Área: Anatomia Patológica

Trabalho efetuado sob a Orientação de:

Doutora Raquel Maria da Silva Graça Almeida

E sob a Coorientação de:

Doutora Maria de Fátima Machado Henriques Carneiro

Trabalho organizado de acordo com as normas da revista:

European Journal of Cancer

março, 2014

FMUP

Eu, Mónica Sofia Gonçalves Garrido, abaixo assinado, nº mecanográfico 200801271, estudante do 6º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 07/03/2014

Assinatura conforme cartão de identificação:

Mónica Sofia Gonçalves Garrido

Projecto de Opção do 6º ano – DECLARAÇÃO DE REPRODUÇÃO

NOME

Mónica Sofia Gonçalves Garrido

CARTÃO DE CIDADÃO

E-MAIL

TELEFONE OU TELEMÓVEL

13780257 9ZZ7

monikikita_garrido@hotmail.com

913131252

NÚMERO DE ESTUDANTE

DATA DE CONCLUSÃO

200801271

2014

DESIGNAÇÃO DA ÁREA DO PROJECTO

Anatomia Patológica

TÍTULO DISSERTAÇÃO

SOX2 and CDX2 in gastric intestinal metaplasia and dysplasia: impact on differentiation

ORIENTADOR

Raquel Maria da Silva Graça Almeida

COORDINADOR

Maria de Fátima Machado Henriques Carneiro

É autorizada a reprodução integral desta Dissertação para efeitos de investigação e de divulgação pedagógica, em programas e projectos coordenados pela FMUP.

Faculdade de Medicina da Universidade do Porto, 07/03/2014

Assinatura conforme cartão de identificação:

Mónica Sofia Gonçalves Garrido

AGRADECIMENTOS

À Professora Doutora Fátima Carneiro, com quem partilhamos mais do que noites de trabalho e dívidas de kitkats. Obrigada por toda a ajuda e dedicação, tantas vezes a horas avançadas da madrugada.

À Doutora Raquel Almeida, por me receber na sua equipa e ter aceite orientar esta tese de mestrado. Obrigada pela disponibilidade, rapidez, eficiência e simplicidade.

À Vânia, por me ter aceite como parceira do seu trabalho. Obrigada pelas gargalhadas e boa disposição que tanto ajudaram a encurtar as horas de trabalho laboratorial.

Ao Franclim, por nos apresentar um pouco do mundo da ciência. Obrigada pelas horas de paciência sempre com boa disposição.

A todo o Serviço de Anatomia Patológica do CHSJ, por acudirem a todos os nossos pedidos de ajuda. Obrigada por nos acolherem na sua casa.

À Irene, que nos contagiou com a sua amizade e paixão pela patologia. Obrigada por nos acompanhar de perto nesta jornada.

À família, pelas longas horas de ausência. Obrigada por tudo compreenderem e apoiarem.

Aos amigos, a quem não é possível tanto agradecer.

Ao Pedro por me receber na sua vida. Obrigada por estar presente em cada minuto desta jornada.

À nossa festa.

INTRODUCTION

Gastric Cancer – Overview

Gastric cancer was the most common and most lethal cancer in the world during most part of the last century. Despite a gradual decrease in its incidence, it is still the fifthmost common malignancy (6.8% of all cancers), after lung, breast, colo-rectum and prostate and the third leading cause of cancer death (8.8% of total) after lung and liver cancers. More than 70% of the cases occur in developing countries.¹

In Portugal, the scenario is similar. Despite a slow decrease in stomach cancer mortality, it still represents one of the most important oncological problems.² According to IARC Cancer Base, gastric cancer is the fifth most incident cancer in Portugal (6.1% of all cancers) and the third most deadly (9.5% of total). These values surpass the ones of WHO Europe Region (4.3% and 6.5% respectively).^{1, 3}

More than 90% of gastric tumours are adenocarcinomas.⁴ According to Lauren's classification, these can be further divided into two main histopathological entities: intestinal and diffuse types.⁵ They differ in epidemiologic, clinic, pathologic and molecular patterns.⁶

Intestinal type gastric carcinoma is the most common subtype and it is preceded by a prolonged precancerous cascade, most frequently initiated by *Helicobacter pylori*, that is still not fully understood.^{6, 7} Morphologically, it forms glands that range from well differentiated to moderately differentiated tumours and it typically arises on a background of chronic atrophic gastritis with intestinal metaplasia.⁶

Diffuse type gastric carcinoma is not preceded by well-defined precancerous lesions.⁷ It also shows an association with *H. pylori* but, unlike intestinal type, it develops without passing through the intermediate step of intestinal metaplasia.^{8, 9} It may occur in an hereditary context, but most frequently it is sporadic. In both cases, loss of expression

of the cellular adhesion protein E-cadherin results, morphologically, in poorly cohesive cells diffusely infiltrating the gastric wall with little or no gland formation.⁶

Gastric Cancer – Carcinogenesis Pathways

Nowadays, the most accepted model of gastric carcinogenesis of the intestinal type still has its foundations on Correa's model. According to Correa, a decades-long, multistep and multifactorial cascade starting with chronic gastric inflammation, progressing to gland loss and their replacement by intestinal-type epithelium, followed by loss of cellular differentiation, leads ultimately to the development of invasive gastric adenocarcinoma (figure 1).^{7, 10-12}




Figure 1 - Correa's Carcinogenesis Model

Tsukashita et al. have suggested that the tumourigenesis of low-grade dysplasia particularly with an intestinal phenotype may be different from high-grade dysplasia and intramucosal carcinomas with gastric phenotype.¹³ Nishimura et al. tested this hypothesis and concluded that non-invasive low-grade neoplasms of the intestinal lineage progress to non-invasive high-grade neoplasms but rarely to intramucosal adenocarcinomas, whereas intramucosal adenocarcinomas of the gastric lineage largely arise *de novo* from the proper gastric mucosa and are partially derived from non-invasive high-grade neoplasms.¹⁴

These two models are not necessarily mutually exclusive so, although gastric carcinogenesis is not fully understood, we can admit that adenoma-carcinoma sequence is responsible for a subset of gastric cancers and that the rest of them may arise *de novo*.

Risk Factors and Determinants of Gastric Carcinogenesis

In its gastric carcinogenesis model, Correa also proposed that both deleterious and protective factors could modulate the progression/regression of this cascade.¹⁰

Helicobacter pylori is the most frequent cause of gastritis, being classified by IARC as a type I (definite) human carcinogen in 1994.¹⁵ Two proposed mechanisms to step in the carcinogenic cascade are the immune response elicited by *H. pylori* and the damage resulting from oxidative stress.¹⁶ The severity of inflammation varies both with the virulence of *H. pylori* strains, given by *vacA* and *cagA* genotypes, and with proinflammatory host genetic polymorphisms in the *IL1 β* , *IL1RN* and *TNF α* genes.¹⁷⁻¹⁹ Eradication with antibiotic treatment can reverse this initial inflammatory reaction but intestinal metaplasia reversibility is still a controversial issue.²⁰ Furthermore, intestinal metaplasia arises in only 30% of infected individuals, from which only around 7% will develop gastric cancer,^{9, 21} so the presence of other genetic/environmental factors modulating progression towards cancer has to be considered.

Other risk factors for the development of sporadic gastric cancer include salt and nitrous compounds of the diet and tobacco smoking. On the other side, fresh fruits and vegetables may play a protective role.²²

Pre-malignant conditions and lesions - metaplasia and dysplasia

Multifocal atrophic gastritis and intestinal metaplasia confer a high risk for the development of gastric cancer, thus they are considered to be precancerous conditions.²³

Intestinal metaplasia represents a phenotypic change from the normal epithelial cell of gastric mucosa to an intestinal phenotype. One of its classifications divides IM into complete or incomplete type, both characterized by the presence of intestinal markers such as the transcription factor CDX2 and the mucin MUC2. Complete intestinal

metaplasia displays goblet and absorptive cells and no expression of gastric mucins. Incomplete intestinal metaplasia displays goblet and columnar non-absorptive cells, in which gastric mucins (MUC5AC, and MUC6) are co-expressed with the intestinal markers.²⁴ This classification has clinical relevance since incomplete intestinal metaplasia confers a higher risk of progression to gastric cancer.²⁵

Gastric dysplasia (also known as intraepithelial neoplasia or non-invasive neoplasia) represents the penultimate stage of the gastric carcinogenesis sequence. Histologically, it is defined by unequivocal neoplastic epithelium without evidence of tissue invasion, and is thus a direct neoplastic precancerous lesion.²³ The term adenoma refers to a raised circumscribed dysplastic lesion protruding above the mucosal surface.²⁶

Observation of different degrees of cellular and architectural atypia allows classification into low and high grade dysplasia.²⁷ Most patients harboring lesions classified as high grade dysplasia are at high risk for either synchronous invasive carcinoma or its rapid development.^{28, 29}

Dysplasia can also be classified, according to its phenotypical characteristics, into adenomatous/intestinal or foveolar/gastric.³⁰ These morphological types can be confirmed by immunohistochemistry techniques. Intestinal type has positivity to MUC2, CD10 and CDX2 and gastric type has positivity to MUC5AC and MUC6, lack of CD10 and low CDX2 expression. Those which show both gastric and intestinal phenotypes are classified as hybrid/mixed phenotype while those showing neither gastric nor intestinal markers are grouped as null type.⁶

Although the literature on the clinical relevance of this morphological classification is limited, some clues have aroused. Park et al have showed that adenomatous type dysplasia is associated with IM showing a complete intestinal phenotype, while foveolar and hybrid dysplasias are closer to incomplete IM.³¹ Also, Tsukashita et al

demonstrated that the majority of low-grade dysplasia (81.8 %) expresses intestinal markers and generally expresses no gastric markers, whereas more than half of high-grade dysplasia expresses gastric markers (72.2%).^{13, 31}

Molecular Pathology

The research on the determinants of gastric cancer precursors has been less extensive than for the cancer endpoints and there are currently no immunohistochemical or molecular assays that can help stratify the risk of progression of gastric dysplastic lesions.

The genetic and molecular abnormalities occurring during gastric carcinogenesis include chromosomal instability, CpG-island methylation and microsatellite instability (MSI).³⁰

Chromosomal instability refers to copy number variations and is the predominant type of genomic instability, being present in more than 60% of gastric cancers.^{32, 33} Loss of heterozygosity at the *TP53* locus was demonstrated in 14% of a series of metaplastic lesions, in 22% of gastric dysplastic lesions and 30% to 50% of gastric carcinomas, consistent with their role in cancer progression.³⁰

Aberrant DNA hypermethylation occurs preferentially in CpG islands.³³ Among gastric adenomas, it was described in genes involved in cell cycle regulation (*p14*, *p16*, *COX2*), signal transduction (*APC*), DNA repair (*hMLH1* and *MGMT*) and invasion and metastasis (*E-cadherin* and *TIMP3*).³⁰ *RUNX3* is a recently recognized tumor suppressor gene for gastric cancer. An increase in the proportion of its promoter methylation along gastric carcinogenesis was described, with 16% in chronic atrophic gastritis, 37% in IM, 42% in gastric adenoma, 55% in dysplasia, and 75% in GC tissues, suggesting its importance at the later stages of gastric carcinogenesis.^{30, 33}

MSI phenotype results from accumulating genomic mutations due to inactivation of the DNA mismatch repair pathway, through promoter hypermethylation and loss of expression of *MHL1*. MSI has been documented both in intestinal metaplasia and in dysplastic lesions (~20%) being therefore considered among the early molecular events in gastric carcinogenesis.³²

Based on gene expression profiling, three gastric tumor subtypes were identified. They were classified as proliferative, mesenchymal, and metabolic and they show differential sensitivities to chemotherapy agents.³⁴

Although molecular events underlying gastric carcinogenesis are not fully uncovered, it is becoming evident that the balance between gastric and intestinal differentiation has a significant impact, at least in the early steps of the gastric carcinogenesis cascade, with the most frequent pre-malignant condition resulting from a switch of gastric to an intestinal differentiation profile.

CDX2 – An intestinal differentiation transcription factor

CDX2 is a transcription factor member of the caudal-related homeobox gene family with an important role for intestinal epithelial development.³⁵ In normal conditions, *CDX2* is expressed in small intestinal and colonic epithelia, but not in gastric epithelium.³⁶ However, in certain unfavourable conditions to the normal gastric mucosa environment – usually elicited by *H. pylori* infection – gastric epithelial cells may acquire ectopic expression of *CDX2* and transdifferentiate into an intestinal phenotype – intestinal metaplasia.³⁷⁻⁴⁰

The strength of these observations largely come from transgenic mouse models studies, in which *CDX2* was under the control of promoters from different gastric-specific genes (*Foxa3* or *H+/K+-ATPase b-subunit*). Mutoh et al. and Silberg et al. reported that the ectopic expression of *CDX2* in the gastric mucosa of transgenic mice alone was sufficient to induce a complete transformation of the gastric mucosal glands

to intestinal-like mucosa, confirming the crucial role of CDX2 in intestinal differentiation.^{37, 38} A posterior paper of Mutoh et al. also showed that long-term intestinal metaplasia of the CDX2 transgenic mice induced invasive gastric carcinoma.⁴¹

Some degree of this CDX2 abnormal expression is maintained in the progression to gastric dysplasia and gastric cancers.^{39, 42} Although Kim et al demonstrated a positive correlation between CDX2 expression and the increasing grade of dysplasia and carcinoma,⁴³ more recent papers described the opposite association, with CDX2 expression progressively reduced in gastric dysplasia and cancer.^{42, 44, 45} Other authors reported a significantly decreased CDX2 expression in incomplete intestinal metaplasia compared with complete intestinal metaplasia,^{39, 44} and an inverse association between CDX2 expression in gastric cancer and the expression of gastric mucins MUC5AC and MUC6.^{42, 44} Also, CDX2-positive tumours have a better outcome than CDX2-negative tumours, with less invasiveness, fewer lymph node metastases and a higher 5-year survival.^{42, 45} Based on these findings, a tumour suppressor role was suggested for CDX2 in human gastric carcinogenesis,^{42, 45} similarly to colorectal cancer.^{46, 47}

SOX2 – A gastric differentiation transcription factor

SOX proteins are important transcription factors with pleiotropic functions – stem and progenitor cell fate determination, cell differentiation, proliferation, reprogramming, tissue homeostasis and repair.⁴⁸ They constitute a large family of genes – with 20 different SOX genes identified until now⁴⁹ – and they have in common the HMG domain similarity with Sry (sex-determining region Y).⁴⁸

In normal tissues, SOX2 is expressed in the foregut-derived organs such as pharynx, oesophagus and stomach, but not in hindgut-derived organs such as intestine, where the CDX1 and CDX2 proteins assume the responsibility of cell differentiation.^{50, 51}

Several other tissues also have been shown to express SOX2, among them are trachea, lung, tongue, cervix, brain, skin and bone.⁴⁸

In normal stomach, SOX2 has a nuclear expression, mainly in the neck region of the gastric glands, both at the body and antrum regions. Starting with *H. pylori* stimuli,⁴⁰ a progressive reduction of the expression of SOX2 from normal gastric mucosa to incomplete intestinal metaplasia, then to complete intestinal metaplasia⁵¹ and to adenocarcinoma,^{50, 52} has been demonstrated, along with an inverse increase of CDX2 expression.⁵¹ This *CDX2*-SOX2 interplay was comprehensively demonstrated both with in vitro^{40, 53, 54} and with transgenic mouse models studies. Conditional knock-out of the *CDX2* gene in mice showed increased SOX2 expression in the regions where CDX2 expression was deficient, with a consequent shift from an intestinal to a gastric/esophageal differentiation.^{55, 56} The same was observed when ectopic expression of SOX2 was induced in the intestine.⁵⁷ Also, it has been shown that SOX2 up-regulates stomach-specific expression of *pepsinogen A*⁵⁸ and *MUC5AC* genes.⁵⁹ Another finding is the higher SOX2 expression in adenocarcinomas expressing gastric mucins compared to adenocarcinomas expressing intestinal mucins.^{50, 60} For all these reasons, SOX2 is a highly suspected candidate for gastric differentiation.

SOX2 expression has been positively associated with tumour grade and with worse prognosis in a variety of tumours.⁶¹ Concerning gastric adenocarcinoma, conflicting results were published. Matsuoka et al found a statistically significant correlation between SOX2-positive expression and depth of tumour invasion, lymph node metastasis and lymphatic invasion.⁶² On the other hand, Otsubo et al reported that SOX2 expression is frequently down-regulated in human gastric cancer, through aberrant DNA methylation, associating with a significantly shorter survival time. Also, they showed that SOX2 inhibits proliferation, induces cell-cycle arrest and apoptosis in gastric cell lines, suggesting a tumour suppressor role for SOX2 in gastric carcinogenesis.⁵²

Aim

This work aims to better characterize the balance of gastric vs. intestinal differentiation in the precancerous conditions and lesions of the gastric carcinogenesis cascade through a systematic description of SOX2 expression in the normal gastric mucosa, complete IM, incomplete IM and dysplasia.

References

1. Ferlay J, Soerjomataram I, Ervik M, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F. GLOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 11 [Internet]. Lyon, France: International Agency for Research on Cancer; 2013. Available from: <http://globocan.iarc.fr>, accessed on 04/02/2014.
2. Pinheiro PS, Tyczyński JE, Bray F, *et al.* Cancer incidence and mortality in Portugal. *European journal of cancer* 2003;**39**(17):2507-2520.
3. Carrilho MJ, Patrício L. The Demographic Changes in Portugal. *Revista de Estudos Demográficos* 2010;**48**:44.
4. Crew KD, Neugut AI. Epidemiology of gastric cancer. *World J Gastroenterol* 2006;**12**(3):354-62.
5. Lauren P. The two histological main types of gastric carcinoma: diffuse and so-called intestinal-type carcinoma. An attempt at a histo-clinical classification. *Acta Pathol Microbiol Scand* 1965;**64**:31-49.
6. Bosman FT, Carneiro F, Hruban RH, *et al.* WHO Classification of Tumours of the Digestive System. 4th ed. Lyon: IARC; 2010.
7. Correa P. A Human Model of Gastric Carcinogenesis. *Cancer Research* 1988;**48**(13):3554-3560.
8. Marcos-Pinto R, Dinis-Ribeiro M, Carneiro F, *et al.* First-degree relatives of early-onset gastric cancer patients show a high risk for gastric cancer: phenotype and genotype profile. *Virchows Arch* 2013;**463**(3):391-9.
9. Uemura N, Okamoto S, Yamamoto S, *et al.* Helicobacter pylori infection and the development of gastric cancer. *N Engl J Med* 2001;**345**(11):784-9.
10. Correa P. Human Gastric Carcinogenesis: A Multistep and Multifactorial Process—First American Cancer Society Award Lecture on Cancer Epidemiology and Prevention. *Cancer Research* 1992;**52**(24):6735-6740.

11. Correa P, Haenszel W, Cuello C, *et al.* A Model for Gastric Cancer Epidemiology. *The Lancet* 1975;**306**(7924):58-60.
12. Pelayo C, Piazzuelo MB. The gastric precancerous cascade. *Journal of Digestive Diseases* 2012;**13**(1):2-9.
13. Tsukashita S, Kushima R, Bamba M, *et al.* MUC gene expression and histogenesis of adenocarcinoma of the stomach. *Int J Cancer* 2001;**94**(2):166-70.
14. Nishimura R, Mukaisho KI, Yamamoto H, *et al.* Precursor-derived versus de-novo carcinogenesis depends on lineage-specific mucin phenotypes of intramucosal gland-forming gastric neoplasms. *Histopathology* 2013.
15. Møller H, Heseltine E, Vainio H. Working group report on schistosomes, liver flukes and *Helicobacter pylori*. *International Journal of Cancer* 1995;**60**(5):587-589.
16. Correa P, Schneider BG. Etiology of Gastric Cancer: What Is New? *Cancer Epidemiology Biomarkers & Prevention* 2005;**14**(8):1865-1868.
17. Machado JC, Figueiredo C, Canedo P, *et al.* A proinflammatory genetic profile increases the risk for chronic atrophic gastritis and gastric carcinoma. *Gastroenterology* 2003;**125**(2):364-71.
18. Xue H, Lin B, Ni P, *et al.* Interleukin-1B and interleukin-1 RN polymorphisms and gastric carcinoma risk: a meta-analysis. *J Gastroenterol Hepatol* 2010;**25**(10):1604-17.
19. Costa AC, Figueiredo C, Touati E. Pathogenesis of *Helicobacter pylori* infection. *Helicobacter* 2009;**14 Suppl 1**:15-20.
20. Barros R, Freund J-N, David L, *et al.* Gastric intestinal metaplasia revisited: function and regulation of CDX2. *Trends in molecular medicine* 2012;**18**(9):555-563.
21. Hsu PI, Lai KH, Hsu PN, *et al.* *Helicobacter pylori* infection and the risk of gastric malignancy. *Am J Gastroenterol* 2007;**102**(4):725-30.
22. Peleteiro Br, Lunet N. Role of Genetic and Environmental Risk Factors in Gastric Carcinogenesis Pathway. In: Tonino DP, editor. Gastritis and Gastric Cancer - New Insights in Gastroprotection, Diagnosis and Treatments: InTech; 2011.

23. Dinis-Ribeiro M, Areia M, Vries AC, *et al.* Management of precancerous conditions and lesions in the stomach (MAPS): guideline from the European Society of Gastrointestinal Endoscopy (ESGE), European Helicobacter Study Group (EHSO), European Society of Pathology (ESP), and the Sociedade Portuguesa de Endoscopia Digestiva (SPED). *Virchows Archiv* 2012;**460**(1):19-46.
24. Silva E, Teixeira A, David L, *et al.* Mucins as key molecules for the classification of intestinal metaplasia of the stomach. *Virchows Arch* 2002;**440**(3):311-7.
25. González CA, Sanz-Anquela JM, Gisbert JP, *et al.* Utility of subtyping intestinal metaplasia as marker of gastric cancer risk. A review of the evidence. *International Journal of Cancer* 2013;**133**(5):1023-1032.
26. Lauwers GY, Riddell RH. Gastric epithelial dysplasia. *Gut* 1999;**45**(5):784-90.
27. Kelly PJ, Lauwers GY. Gastric polyps and dysplasia. *Diagnostic Histopathology* 2011;**17**(2):50-61.
28. Rugge M, Farinati F, Baffa R, *et al.* Gastric epithelial dysplasia in the natural history of gastric cancer: a multicenter prospective follow-up study. Interdisciplinary Group on Gastric Epithelial Dysplasia. *Gastroenterology* 1994;**107**(5):1288-96.
29. Rugge M, Cassaro M, Di Mario F, *et al.* The long term outcome of gastric non-invasive neoplasia. *Gut* 2003;**52**(8):1111-6.
30. Tan D, Lauwers G. *Advances in Surgical Pathology: Gastric Cancer*: Lippincott Williams & Wilkins - Wolters Kluwer; 2011.
31. Park do Y, Srivastava A, Kim GH, *et al.* Adenomatous and foveolar gastric dysplasia: distinct patterns of mucin expression and background intestinal metaplasia. *Am J Surg Pathol* 2008;**32**(4):524-33.
32. Fassan M, Baffa R, Kiss A. Advanced precancerous lesions within the GI tract: the molecular background. *Best Pract Res Clin Gastroenterol* 2013;**27**(2):159-69.
33. Figueiredo C, Garcia-Gonzalez MA, Machado JC. Molecular pathogenesis of gastric cancer. *Helicobacter* 2013;**18 Suppl 1**:28-33.

34. Lei Z, Tan IB, Das K, *et al.* Identification of molecular subtypes of gastric cancer with different responses to PI3-kinase inhibitors and 5-fluorouracil. *Gastroenterology* 2013;**145**(3):554-65.
35. Guo RJ, Suh ER, Lynch JP. The role of Cdx proteins in intestinal development and cancer. *Cancer Biol Ther* 2004;**3**(7):593-601.
36. Yuasa Y. Control of gut differentiation and intestinal-type gastric carcinogenesis. *Nat Rev Cancer* 2003;**3**(8):592-600.
37. Silberg DG, Sullivan J, Kang E, *et al.* Cdx2 ectopic expression induces gastric intestinal metaplasia in transgenic mice. *Gastroenterology* 2002;**122**(3):689-96.
38. Mutoh H, Hakamata Y, Sato K, *et al.* Conversion of gastric mucosa to intestinal metaplasia in Cdx2-expressing transgenic mice. *Biochem Biophys Res Commun* 2002;**294**(2):470-9.
39. Almeida R, Silva E, Santos-Silva F, *et al.* Expression of intestine-specific transcription factors, CDX1 and CDX2, in intestinal metaplasia and gastric carcinomas. *J Pathol* 2003;**199**(1):36-40.
40. Camilo V, Barros R, Sousa S, *et al.* Helicobacter pylori and the BMP pathway regulate CDX2 and SOX2 expression in gastric cells. *Carcinogenesis* 2012;**33**(10):1985-92.
41. Mutoh H, Sakurai S, Satoh K, *et al.* Development of gastric carcinoma from intestinal metaplasia in Cdx2-transgenic mice. *Cancer Res* 2004;**64**(21):7740-7.
42. Park do Y, Srivastava A, Kim GH, *et al.* CDX2 expression in the intestinal-type gastric epithelial neoplasia: frequency and significance. *Mod Pathol* 2010;**23**(1):54-61.
43. Kim HS, Lee JS, Freund JN, *et al.* CDX-2 homeobox gene expression in human gastric carcinoma and precursor lesions. *J Gastroenterol Hepatol* 2006;**21**(2):438-42.
44. Liu Q, Teh M, Ito K, *et al.* CDX2 expression is progressively decreased in human gastric intestinal metaplasia, dysplasia and cancer. *Mod Pathol* 2007;**20**(12):1286-97.

45. Qin R, Wang NN, Chu J, *et al.* Expression and significance of homeodomain protein Cdx2 in gastric carcinoma and precancerous lesions. *World J Gastroenterol* 2012;**18**(25):3296-302.
46. Mallo GV, Rechreche H, Frigerio JM, *et al.* Molecular cloning, sequencing and expression of the mRNA encoding human Cdx1 and Cdx2 homeobox. Down-regulation of Cdx1 and Cdx2 mRNA expression during colorectal carcinogenesis. *Int J Cancer* 1997;**74**(1):35-44.
47. Choi BJ, Kim CJ, Cho YG, *et al.* Altered expression of CDX2 in colorectal cancers. *Apmis* 2006;**114**(1):50-4.
48. Sarkar A, Hochedlinger K. The Sox Family of Transcription Factors: Versatile Regulators of Stem and Progenitor Cell Fate. *Cell Stem Cell* 2013;**12**(1):15-30.
49. Schepers GE, Teasdale RD, Koopman P. Twenty pairs of sox: extent, homology, and nomenclature of the mouse and human sox transcription factor gene families. *Dev Cell* 2002;**3**(2):167-70.
50. Li XL, Eishi Y, Bai YQ, *et al.* Expression of the SRY-related HMG box protein SOX2 in human gastric carcinoma. *Int J Oncol* 2004;**24**(2):257-63.
51. Tsukamoto T, Inada K, Tanaka H, *et al.* Down-regulation of a gastric transcription factor, Sox2, and ectopic expression of intestinal homeobox genes, Cdx1 and Cdx2: inverse correlation during progression from gastric/intestinal-mixed to complete intestinal metaplasia. *J Cancer Res Clin Oncol* 2004;**130**(3):135-45.
52. Otsubo T, Akiyama Y, Yanagihara K, *et al.* SOX2 is frequently downregulated in gastric cancers and inhibits cell growth through cell-cycle arrest and apoptosis. *Br J Cancer* 2008;**98**(4):824-31.
53. Benahmed F, Gross I, Gaunt SJ, *et al.* Multiple regulatory regions control the complex expression pattern of the mouse Cdx2 homeobox gene. *Gastroenterology* 2008;**135**(4):1238-1247, 1247.e1-3.
54. Asonuma S, Imatani A, Asano N, *et al.* Helicobacter pylori induces gastric mucosal intestinal metaplasia through the inhibition of interleukin-4-mediated HMG box

protein Sox2 expression. *Am J Physiol Gastrointest Liver Physiol* 2009;**297**(2):G312-22.

55. Stringer EJ, Duluc I, Saandi T, *et al.* Cdx2 determines the fate of postnatal intestinal endoderm. *Development* 2012;**139**(3):465-74.

56. Gao N, White P, Kaestner KH. Establishment of intestinal identity and epithelial-mesenchymal signaling by Cdx2. *Dev Cell* 2009;**16**(4):588-99.

57. Raghoebir L, Bakker ER, Mills JC, *et al.* SOX2 redirects the developmental fate of the intestinal epithelium toward a premature gastric phenotype. *J Mol Cell Biol* 2012;**4**(6):377-85.

58. Tani Y, Akiyama Y, Fukamachi H, *et al.* Transcription factor SOX2 up-regulates stomach-specific pepsinogen A gene expression. *J Cancer Res Clin Oncol* 2007;**133**(4):263-9.

59. Park ET, Gum JR, Kakar S, *et al.* Aberrant expression of SOX2 upregulates MUC5AC gastric foveolar mucin in mucinous cancers of the colorectum and related lesions. *Int J Cancer* 2008;**122**(6):1253-60.

60. Tsukamoto T, Mizoshita T, Mihara M, *et al.* Sox2 expression in human stomach adenocarcinomas with gastric and gastric-and-intestinal-mixed phenotypes. *Histopathology* 2005;**46**(6):649-58.

61. Schoenhals M, Kassambara A, Vos JD, *et al.* Embryonic stem cell markers expression in cancers. *Biochemical and Biophysical Research Communications* 2009;**383**(2):157-162.

62. Matsuoka J, Yashiro M, Sakurai K, *et al.* Role of the Stemness Factors Sox2, Oct3/4, and Nanog in Gastric Carcinoma. *Journal of Surgical Research* 2012;**174**(1):130-135.

Differentiation reprogramming in gastric intestinal metaplasia and dysplasia: role of SOX2 and CDX2

Vânia Camilo^a, Mónica Garrido^b, Pedro Valente^b, Sara Ricardo^a, Sofia Sousa^a, Ana Luísa Amaral^a, Rita Barros^a, Paula Chaves^c, Fátima Carneiro^{a,b,d}, Leonor David^{a,b}, Raquel Almeida^{a,b}

^aInstitute of Molecular Pathology and Immunology of the University of Porto (IPATIMUP), Porto, Portugal

^bFaculty of Medicine of the University of Porto, Porto, Portugal

^cPortuguese Oncology Institute Francisco Gentil, E.P.E., Lisbon, Portugal

^dPathology Department, Centro Hospitalar S. João, Porto, Portugal

Corresponding author: Raquel Almeida, IPATIMUP, Rua Dr. Roberto Frias s/n, 4200-465 Porto, Portugal. Email: ralmeida@ipatimup.pt; Tele: +351225570700; Fax: +351225570799

ACKNOWLEDGMENTS

This work was supported by “Project NORTE - 07-0124-FEDER-000024 co-financed by Programa Operacional Regional do Norte (ON.2 – O Novo Norte), under Quadro de Referência Estratégico Nacional (QREN), by Fundo Europeu de Desenvolvimento Regional (FEDER).” IPATIMUP is an Associate Laboratory of the Portuguese Ministry of Science, Technology and Higher Education and is partially supported by FCT. V.C. and R.B. acknowledge FCT for financial support (grant numbers SFRH/BD/63300/2009 and SFRH/BPD/68276/2010, respectively).

ABSTRACT

Gastric cancer remains a major health concern worldwide. Intestinal metaplasia (IM), resulting from *de novo* expression of CDX2, and dysplasia are precursor lesions of gastric cancer associated with increased risk for cancer development. Multiple evidences suggest that SOX2 is a transcription factor with a role in gastric differentiation and its interplay with CDX2 may determine gastric carcinogenesis. Our aim was to assess SOX2 involvement in gastric premalignant conditions and lesions in an attempt to establish its relationship with CDX2 and with the differentiation reprogramming that characterizes gastric carcinogenesis. Characterization of gastric (SOX2, MUC5AC, MUC6) and intestinal (CDX2 and MUC2) markers in normal gastric mucosa, in 55 foci of IM and in 26 foci of dysplasia was performed by immunohistochemistry. *In vitro* models were used to study the putative cross-regulation between SOX2 and CDX2. SOX2 was expressed in the normal gastric mucosa, in the presumptive stem cell compartment, and was maintained in 7% of the complete and 85% of the incomplete IM subtypes, the latter defined by MUC5AC expression. 12% of the dysplastic lesions expressed SOX2 and the association with MUC5AC was lost. Conversely, CDX2 was present in all IM and dysplastic lesions. Finally, SOX2 negatively regulated CDX2 in a cell-type specific manner. In conclusion, SOX2 associates with gastric differentiation and is lost in the progression to dysplasia, whereas CDX2 is *de novo* acquired in IM and maintained in dysplasia. This suggests that the balance between gastric and intestinal differentiation programs impacts on progression of the gastric carcinogenic cascade.

Keywords: SOX2, CDX2, intestinal metaplasia, dysplasia, gastric cancer

INTRODUCTION

Gastric cancer is the fourth most common malignancy and the second-leading cause of cancer-related deaths worldwide.¹ The majority of gastric cancers is diagnosed late, and as such has a dismal prognosis, which is reflected by a 5-year survival rate of no more than 25%.² Thus, the most promising strategies to control the disease are based on prevention and early diagnosis. Gastric cancer is usually initiated by an inflammatory process associated with infection by *Helicobacter pylori*, which may lead to multifocal atrophic gastritis, intestinal metaplasia (IM), dysplasia and finally cancer.³ IM is a switch from a gastric to an intestinal differentiation profile and is the most frequent premalignant condition of the stomach, appearing adjacent to more than 80% of gastric cancers.⁴ It is an heterogeneous lesion with two subtypes, complete and incomplete, both characterized by the presence of intestinal markers such as the transcription factor CDX2 and the mucin MUC2. The complete IM subtype is defined by the absence of gastric differentiation markers, such as mucins MUC5AC and MUC6, whereas in the incomplete IM subtype there is concomitant expression of gastric and intestinal markers.⁵ This classification has clinical implications because the incomplete subtype is more frequently associated with gastric cancer and shows an increased risk for cancer progression.^{4,6}

CDX2 is a homeobox transcription factor critical for intestinal differentiation⁷⁻⁸ and is a specific biomarker of the early steps of the gastric carcinogenic cascade, driving the onset of IM.⁹⁻¹⁰ The key role of CDX2 in the metaplastic transformation of the gastric mucosa was categorically demonstrated by two transgenic mouse models with ectopic expression of CDX2 in the gastric epithelium and subsequent development of IM with absorptive, goblet, and enteroendocrine cell types.¹¹⁻¹²

On the other hand, an increasing number of evidences support the involvement of the transcription factor SOX2 in gastric differentiation. SOX2 is a sex-determining region Y-box 2 gene, a member of the high mobility group (HMG) domain proteins that

is essential to maintain pluripotency in embryonic stem (ES) cells and also to reprogram fibroblasts into induced pluripotent (iPS) cells.¹³⁻¹⁴ More recently, it has also been identified as an adult stem cell marker in mice.¹⁵ Furthermore, modulating SOX2 expression in mice demonstrated the importance of this protein in the differentiation of the esophageal epithelium and proper morphogenesis of the esophagus, trachea and lung.¹⁶⁻¹⁷ In the digestive tract, it was observed in chick and in mouse gut that SOX2 and CDX2 expression are mutually exclusive.¹⁸⁻¹⁹ *In vitro* studies showed that SOX2 negatively regulated the CDX2 promoter by hampering the action of other transcription factors in an intestinal context and that SOX2 downregulation led to CDX2 overexpression in a gastric context.²⁰⁻²¹ In accordance with these observations, we have demonstrated that SOX2 and CDX2 can be inversely regulated by the BMP pathway and *Helicobacter pylori*.²² Moreover, two mouse models with conditional CDX2 loss of function, showed increased SOX2 expression in the regions where CDX2 was abrogated, with a consequent shift from an intestinal to a gastric/esophageal differentiation.²³⁻²⁴ The same was observed when forced expression of SOX2 was induced in the intestine,²⁵⁻²⁶ suggesting that the interplay of these two transcription factors is critical for the balance between gastric and intestinal differentiation.

Here, we sought to characterize SOX2 expression and the relationship with CDX2 in gastric cancer precursor lesions in an attempt to establish its involvement in the differentiation reprogramming that characterizes gastric carcinogenesis.

MATERIAL AND METHODS

Material

Normal gastric mucosa was studied in 10 biopsies representative of the body (n = 5) and the antrum (n = 5) regions obtained from individuals with no gastric pathologies. IM (55 foci) and the adjacent gastric mucosa were studied in 11 surgical specimens obtained from patients with gastric carcinoma undergoing surgery at Centro Hospitalar S. João, Porto, Portugal. IM foci were classified into complete (n = 29) or incomplete (n = 26) type according to the pattern of mucin expression.⁵ Twenty-six foci of dysplasia were studied in 22 samples obtained by Endoscopic Submucosal Dissection (ESD) at the same hospital. Foci were considered distinct when they were separated by a stretch of gastric glands. The use of retrospective samples from which informed consent cannot be obtained is authorized for research studies by the Portuguese law.

Immunohistochemistry

Paraffin-embedded specimens were subjected to immunohistochemistry for SOX2, CDX2, MUC5AC, MUC6 and MUC2 using the antibodies described in Table 1. Detection of CDX2, MUC5AC and MUC2 was performed by incubation with a biotin-labeled rabbit anti-mouse secondary antibody (DAKO, 1:100) followed by incubation with an avidin/biotin detection system (Vectastain ABC kit, Vector Laboratories, Burlingame, CA, USA) and development with 3,3'-diaminobenzidine (DAB). Detection of SOX2 was done using the Dako REAL™ Envision™ Detection System Peroxidase/DAB+ (DAKO, Glostrup, Denmark) according to the manufacturer's instructions.

For double staining (SOX2/MUC5AC or SOX2/MUC6), immunohistochemistry was first performed for SOX2 using the Dako REAL™ Envision™ Detection System

Peroxidase/DAB+ (DAKO, Glostrup, Denmark) and immediately followed by immunohistochemistry for MUC5AC or MUC6 using the Envision™ G2 System/ AP (Permanent Red) (DAKO), according to the manufacturer's instructions. Tissue counterstaining was performed using Mayer hematoxylin. Cases were considered positive when more than 5% of the cells were stained with each antibody.

Cell Culture

The human gastric carcinoma cell lines AGS (ATCC) and MKN45 (JCRB0254, The RIKEN Cell Bank) were maintained in RPMI1640-GlutaMAx (Gibco, Invitrogen), whereas the human colon carcinoma cell line Caco-2 (ATCC) was maintained in DMEM (Gibco, Invitrogen). Media were supplemented with 10% fetal bovine serum (Gibco, Invitrogen, Carlsbad, USA) and 1% antibiotics (10U/mL penicillin and 10µg/mL streptomycin) (Gibco, Invitrogen, Carlsbad, USA). Cells were maintained at 37°C in a humidified 5% CO₂ incubator.

Transient transfection assays

For AGS and MKN45, one day prior to transfection, cells (7.5×10^5) were seeded in 6-well plates. Transfection of the expression vector CMV/SOX2, (1µg/well) or the empty vector was carried out using Lipofectamine 2000 (Invitrogen) (1µg DNA:1.5µl lipofectamine ratio) in serum and antibiotic free OPTI-MEM (Gibco, Invitrogen). Cells were recovered 48h after transfection. Caco-2 cells were subjected to two consecutive rounds of transfection. Cells were seeded in 6-well plates and transfected either with the expression vector CMV/SOX2 or with the empty vector (1µg/well), as described above. The medium was changed and a second similar transfection was performed 72 hours after plating. Cells were recovered 48h after the second transfection.

Protein Extraction and Western Blot

Whole cell extracts were obtained by resuspension of cell pellets in RIPA buffer (50mM TrisHCl pH 7.4, 150mM NaCl, 2mM EDTA, 1% NP-40, 0.1% SDS) in the presence of Complete protease inhibitors cocktail, (Roche, Indianapolis, USA) and quantification of total protein was determined by BCA protein assay (Pierce, Illinois, USA). Protein extracts were then subjected to standard SDS-PAGE, transferred to a nitrocellulose membrane (Amersham, GE Healthcare, UK) and blotted with primary antibodies (anti-CDX2, 1:500, Biogenex; anti-SOX2, 1:4500, SIGMA; anti- β -actin, 1:8000, SCBT) in 5% BSA (Sigma, St. Louis, USA) in TBS 0.01% Tween-20 (Sigma, St. Louis, USA). Peroxidase-conjugated secondary antibodies were used and developed with the ECL detection kit (Amersham, GE Healthcare, UK). Quantification of the western blots was performed using the Software Quantity One (BioRad, CA, USA).

RNA extraction and Real-Time PCR

Total RNA was extracted using TRI Reagent (Sigma, St. Louis, USA) and converted to cDNA using the SuperScript® II Reverse Transcriptase (Invitrogen, Carlsbad, USA). CDX2 or 18S were amplified with SYBR Green (Applied Biosystems, Foster City, USA) in a fluorescence reader ABI Prism 7500. The levels of 18S were used for normalization and relative mRNA levels were calculated. Each experiment was carried out in triplicates at least twice; the results are expressed as means \pm SD of representative triplicates.

Statistical analysis

Statistical analysis was performed using the StatView program (version 5.0).

Distributions were compared by χ^2 and Student's *t*-test and significance was assumed whenever *p* values were <0.05.

RESULTS

SOX2 expression in the normal gastric mucosa

We first characterized SOX2 expression in the normal gastric mucosa. For that purpose, we performed immunohistochemistry in 10 normal gastric mucosas and in 11 gastric mucosas adjacent to IM. The pattern of expression was similar in all samples. SOX2 was mainly expressed in the nuclei of the cells in the neck region of the gastric glands, both at the body and antrum (Fig 1) and expression was progressively lost towards the surface. SOX2 expression is also less marked in the deep glands particularly in the body region (Fig 1).

SOX2 expression in gastric premalignant lesions

We next evaluated the expression pattern of SOX2 in gastric premalignant conditions and lesions: intestinal metaplasia and dysplasia. We performed immunohistochemistry in 26 foci of incomplete IM and 29 foci of complete IM, from a total of 21 cases. Complete and incomplete IM were defined, as previously described,⁵ by the expression pattern of gastric mucins, with the complete type characterized by the absence of gastric mucins MUC5AC and MUC6 whereas the incomplete type expresses at least one of these mucins. SOX2 exhibited a distinct expression pattern according to the IM subtype, being expressed in 85% (22/26) of the incomplete IM foci and in 7% (2/29) of the complete IM foci (Table 2). Double immunostainings showed co-expression of SOX2 and MUC5AC in incomplete IM (Fig 2) and absence of both proteins in complete IM (Fig 2). MUC2 and CDX2 were expressed in all IM foci, as previously described (data not shown).⁹

We next evaluated the expression pattern of SOX2 in 26 foci of dysplasia: low grade dysplasia (LGD) – 20 foci; high grade dysplasia – 6 foci. SOX2 was expressed in 12% of the foci (3/26), all of low grade dysplasia (Table 3). Fig 3 shows dysplastic glands with SOX2 expression adjacent to negative ones. To further assess the differentiation characteristics of these dysplastic lesions we studied CDX2, MUC2, MUC5AC and MUC6 expression. The results are displayed in Table 3 and showed that all dysplastic foci expressed CDX2, 46% (12/26) expressed MUC2, 58% (15/26) expressed MUC5AC and 68%(17/25) expressed MUC6. Moreover, CDX2 expression was uniform throughout the lesions, similarly to IM, whereas expression of all mucins was scattered and heterogeneous (Fig 3).

SOX2 and CDX2 cross-regulation

Different studies with mice models suggest that CDX2 negatively regulates SOX2 and also the reverse.^{20-21,25-26} In order to determine if there is a direct cross-regulatory mechanism between SOX2 and CDX2, which could be relevant in the onset of IM, we transiently transfected two gastric carcinoma cell lines (AGS and MKN45) and one colonic carcinoma cell line (Caco-2) with a CMV/SOX2 vector and the gastric carcinoma cell lines with a CMV/CDX2 expression vector (this experiment was only performed in the gastric cell lines since Caco-2 expresses CDX2 at high levels). The results obtained showed that in Caco-2 cells, increased SOX2 levels led to a decreased CDX2 expression both at the mRNA and the protein level (Fig 4) whereas in none of the cell lines CDX2 regulated SOX2 expression (data not shown).

DISCUSSION

We have shown that SOX2 is consistently expressed in the normal gastric mucosa, mostly in the neck region, and in incomplete intestinal metaplasia (IM) conceivably explaining the maintenance of a gastric differentiation program in these cells.

Gastric carcinogenesis is complex and so far largely elusive at the molecular level. However, understanding this process is key in order to design more effective screening and surveillance approaches or even revert preneoplastic lesions. One of the major pathways unfolds in a background of chronic atrophic gastritis and is associated with reprogramming of differentiation leading to IM. It is consensual that IM is an heterogeneous lesion, though different classification criteria have been proposed throughout the years, either based on morphological features or on mucin expression profiles. The mucin expression profile can be determined by classical histochemical methods using the periodic acid-Schiff, Alcian blue, and high iron diamine staining,²⁸ or using immunohistochemistry,⁵ which was the approach applied in this study. Notwithstanding, all the methods lead to the classification of IM into complete and incomplete subtypes, based on loss or persistence of gastric differentiation together with acquisition of the intestinal one as currently accepted by most authors.²⁹ Moreover, incomplete IM, regardless of the classification method used, is consistently more frequently associated with progression to cancer.^{4,6} The involvement of the transcription factor CDX2 in the transformation process leading to both types of IM is undisputable and now an explanation for the gastric phenotype observed in incomplete IM is suggested by the differential expression of SOX2.

Our results clarified the profile of SOX2 expression, both in normal stomach and in gastric lesions, reinforcing its relationship with gastric differentiation.³⁰⁻³¹ We show, for the first time, that SOX2 localizes predominantly to the neck region of the normal gastric epithelium, which is the proliferative compartment of the mucosa and the

expected location of gastric stem cells.³² This observation complies with a role of this transcription factor in the maintenance of the adult gastric epithelium, previously observed in mice¹⁵ and is in accordance with observations that SOX2 regulates MUC5AC and pepsinogen A, both of which are gastric differentiation markers.³³⁻³⁴

Here, we also show that SOX2 is able to repress CDX2 expression, which might be one of the molecular mechanisms involved in maintaining normal gastric differentiation and, when deregulated, in IM onset. In fact, it was shown in other models that SOX2 downregulates CDX2 expression via SOX21.²⁵ This adds to other molecular mechanisms that might also be involved in IM, namely the BMP pathway, that is overexpressed in gastric IM, upregulates CDX2 and downregulates SOX2 in a gastric carcinoma cell line.^{22,35} The potential interactions and or synergies of these different molecular mechanisms need further clarification.

Finally, we disclosed the expression pattern of SOX2 and CDX2 in gastric dysplasia. We showed that the majority of dysplastic lesions do not express SOX2 whereas all of them express CDX2, which is in accordance with Rugge *et al.*³⁶ These two events suggest that the progression from normal gastric mucosa to dysplasia occurs concomitantly with gain of intestinal and loss of gastric differentiation, favoring the theory that dysplasia arises from metaplasia. This complies with the conclusions of Gutierrez-Gonzalez *et al.* showing that IM and dysplasia share genetic mutations in tumor suppressor genes, either APC or p53, in three cases.³⁷ Likewise, Barrett's esophagus which is a very similar lesion to gastric IM, is considered a bona fide preneoplastic lesion where dysplasia ensues. Mutations in p53 or p16 were identified in Barrett's esophagus and dysplasia, further supporting the morphological evidences.³⁸ Our results thus reinforce the link between IM and dysplasia in the gastric setting, based on the expression of SOX2 and CDX2. This observation, strengthened by previous findings that IM is a lesion difficult to revert or even a "point of no return"³⁹ has clinically relevant implications namely regarding surveillance of patients with IM, which

is in line with recent epidemiological data.⁴⁰ Curiously, SOX2 and CDX2 no longer associate with mucin expression in dysplastic foci. A possible interpretation could be that these genes become subjected to regulatory mechanisms associated with the transformation process.

In this study we have demonstrated that SOX2 is expressed in the normal stomach and is maintained in incomplete IM explaining the gastric differentiation still observed in these lesions. Given that SOX2 expression is lost both in dysplasia and in the complete IM subtype, we hypothesize that incomplete IM is a transient state that precedes the onset of both premalignant lesions (Fig 5). In conclusion, the balance between gastric and intestinal differentiation has a significant impact at least in the early steps of gastric carcinogenesis.

Conflict of interest statement

The authors disclose no conflict of interests.

REFERENCES

1. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 2010;**127**:2893-917.
2. Verdecchia A, Francisci S, Brenner H, *et al.* Recent cancer survival in Europe: a 2000-02 period analysis of EURO CARE-4 data. *Lancet Oncol* 2007;**8**:784-96.
3. Correa P. Human gastric carcinogenesis: a multistep and multifactorial process--First American Cancer Society Award Lecture on Cancer Epidemiology and Prevention. *Cancer Res* 1992;**52**:6735-40.
4. Uemura N, Okamoto S, Yamamoto S, *et al.* Helicobacter pylori infection and the development of gastric cancer. *N Engl J Med.* 2001;**345**:784-89.
5. Reis CA, David L, Correa P, Carneiro F, *et al.* Intestinal metaplasia of human stomach displays distinct patterns of mucin (MUC1, MUC2,MUC5AC, and MUC6) expression. *Cancer Res* 1999;**59**:1003-7.
6. González CA, Sanz-Anquela JM, Gisbert JP, Correa P. Utility of subtyping intestinal metaplasia as a marker of gastric cancer risk. A review of the evidence. *Int J Cancer* 2013;**133**:1023-33.
7. Chawengsaksophak K, James R, Hammond VE, Köntgen F, Beck F. Homeosis and intestinal tumours in Cdx2 mutant mice. *Nature* 1997;**386**:84-87.
8. Beck F, Chawengsaksophak K, Waring P, Playford RJ, Furness JB. Reprogramming of the intestinal differentiation and intercalary regeneration in Cdx2 mutant mice. *Proc Natl Acad Sci USA* 1999;**96**:7318-23.
9. Almeida R, Silva E, Santos-Silva F, *et al.* Expression of intestine-specific transcription factors, CDX1 and CDX2, in intestinal metaplasia and gastric carcinomas. *J Pathol* 2003;**199**:36-40.

10. Eda A, Osawa H, Satoh K, *et al.* Aberrant expression of CDX2 in Barrett's epithelium and inflammatory esophageal mucosa. *J Gastroenterol* 2003;**38**:14-22.
11. Mutoh H, Hakamata Y, Sato K, *et al.* Conversion of gastric mucosa to intestinal metaplasia in Cdx2-expressing transgenic mice. *Biochem Biophys Res Commun* 2002;**294**:470-79.
12. Silberg DG, Sullivan J, Kang E, *et al.* Cdx2 ectopic expression induces gastric intestinal metaplasia in transgenic mice. *Gastroenterology* 2002;**122**:689-96.
13. Masui S, Nakatake Y, Toyooka Y, *et al.* Pluripotency governed by Sox2 via regulation of Oct3/4 expression in mouse embryonic stem cells. *Nat Cell Biol* 2007;**9**:625-35.
14. Takahashi K, Yamanaka S. Induction of pluripotency stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 2006;**126**:663-76.
15. Arnold K, Sarkar A, Yram MA, *et al.* SOX2(+) adult stem and progenitor cells are important for tissue regeneration and survival of mice. *Cell Stem Cell* 2011;**9**:317-29.
16. Que J, Okubo T, Goldenring JR, *et al.* Multiple dose-dependent roles for Sox2 in the patterning and differentiation of anterior foregut endoderm. *Development* 2007;**134**:2521-31.
17. Gontan C, de Munck A, Vermeij M, Grosveld F, Tibboel D, Rottier R. Sox2 is important for two crucial processes in lung development: branching morphogenesis and epithelial cell differentiation. *Dev Biol* 2008;**317**:296-309.
18. Ishii Y, Rex M, Scotting PJ, Yasugi S. Region-specific expression of chicken Sox2 in the developing gut and lung epithelium: regulation by epithelial-mesenchymal interactions. *Dev Dyn* 1998;**213**:464-75.
19. Sherwood RI, Chen TY, Melton DA. Transcription dynamics of endodermal organ formation. *Dev Dyn* 2009;**238**:29-42.

20. Benahmed F, Gross I, Gaunt SJ, *et al.* Multiple regulatory regions control the complex expression pattern of the mouse Cdx2 homeobox gene. *Gastroenterology* 2008;**135**:1238-47.
21. Asonuma S, Imatani A, Asano N, *et al.* Helicobacter pylori induces gastric mucosal intestinal metaplasia through the inhibition of interleukin-4-mediated HMG box protein Sox2 expression. *Am J Physiol Gastrointest Liver Physiol* 2009;**297**:312-22.
22. Camilo V, Barros R, Sousa S, *et al.* Helicobacter pylori and the BMP pathway regulate CDX2 and SOX2 expression in gastric cells. *Carcinogenesis* 2012;**33**:1985-92.
23. Gao N, White P, Kaestner KH. Establishment of intestinal identity and epithelial-mesenchymal signaling by Cdx2. *Dev Cell* 2009;**16**:588-99.
24. Stringer EJ, Pritchard CA, Beck F. Cdx2 initiates histodifferentiation of the midgut endoderm. *FEBS Lett* 2008;**582**:2555-60.
25. Kuzmichev AN, Kim SK, D'Alessio AC, *et al.* Sox2 acts through Sox21 to regulate transcription in pluripotent and differentiated cells. *Curr Biol* 2012;**22**:1705-10.
26. Raghoebir L, Bakker ER, Mills JC, *et al.* SOX2 redirects the developmental fate of the intestinal epithelium toward a premature gastric phenotype. *J Mol Cell Biol* 2012;**4**:377-85.
27. Reis CA, David L, Carvalho F, *et al.* Immunohistochemical study of the expression of MUC6 mucin and co-expression of other secreted mucins (MUC5AC and MUC2) in human gastric carcinomas. *J Histochem Cytochem* 2000;**48**:377-88.
28. Jass JR, Filipe MI. A variant of intestinal metaplasia associated with gastric carcinoma: a histochemical study. *Histopathology* 1979;**3**:191-99.
29. Dinis-Ribeiro M, Areia M, de Vries AC, *et al.* Management of precancerous conditions and lesions in the stomach (MAPS):guideline for the European Society of

Gastrointestinal Endoscopy (ESGE), European Helicobacter Study Group (EHSG), European Society of Pathology (ESP), and the Sociedade Portuguesa de Endoscopia digestiva (SPED). *Endoscopy* 2012;**44**: 74–94.

30. Tsukamoto T, Inada K, Tanaka H, *et al.* Down-regulation of a gastric transcription factor, SOX2, and ectopic expression of intestinal homeobox genes, CDX1 and CDX2: inverse correlation during progression from gastric/intestinal-mixed to complete intestinal metaplasia. *J Cancer Res Clin Oncol* 2004;**130**:135-45.

31. Bornschein J, Tóth K, Selgard M, *et al.* Dysregulation of CDX1, CDX2 and SOX2 in patients with gastric cancer also affects the non-malignant mucosa. *J Clin Pathol* 2013;**66**:819-22.

32. Karam SM, Leblond CP. Dynamics of epithelial cells in the corpus of the mouse stomach I. Identification of proliferative cell types and pinpointing of the stem cell. *The Anatomical Record* 1993;**236**:259–79.

33. Park ET, Gum JR, Kakar S, Kwon SW, Deng G, Kim YS. Aberrant expression of SOX2 upregulates MUC5AC gastric foveolar mucin in mucinous cancers of the colorectum and related lesions. *Int J Cancer* 2008;**122**:1253-60.

34. Tani Y, Akiyama Y, Fukamachi H, Yanagihara K, Yuasa Y. Transcription factor SOX2 up-regulates stomach-specific pepsinogen A gene expression. *J Cancer Res Clin Oncol* 2007;**133**:263-69.

35. Barros R, Pereira B, Duluc I, *et al.* Key elements of the BMP/SMAD pathway co-localize with CDX2 in intestinal metaplasia and regulate CDX2 expression in human gastric cell lines. *J Pathol* 2008;**21**:411-20.

36. Rugge M, Ingravallo G, Farinati F, Russo VM, Zaninotto G, Alvisi V. CDX2 homeotic gene expression in gastric noninvasive neoplasia. *Am J Surg Pathol* 2004;**28**:834-35.

37. Gutierrez-Gonzalez L, Graham TA, Rodriguez-Justo M, *et al.* The clonal origins of dysplasia from intestinal metaplasia in the human stomach. *Gastroenterology* 2011;**140**:1251-60.
38. Wang DH, Souza RF. Biology of Barrett's esophagus and esophageal adenocarcinoma. *Gastrointest Endosc Clin N Am* 2011;**21**:25-38.
39. Barros R, da Costa LT, Pinto-de-Sousa J, *et al.* CDX2 autoregulation in human intestinal metaplasia of the stomach: impact on the stability of the phenotype. *Gut* 2011;**60**:290-8.
40. Den Hoed CM, Holster IL, Capelle LG *et al.* Follow-up of premalignant lesions in patients at risk for progression to gastric cancer. *Endoscopy* 2013;**45**:249-56.

FIGURE LEGENDS

Figure 1- SOX2 expression in the normal gastric mucosa. A and C) H&E staining of gastric mucosa from incisura and antrum, respectively. (B and D) Immunodetection of SOX2 (brown) in the same regions of the gastric mucosa.

Figure 2- SOX2 expression in intestinal metaplasia. H&E staining showing gastric mucosa with intestinal metaplasia (A). Double immunodetection of SOX2 (brown) and MUC5AC (red) showing the presence of complete (MUC5AC negative) and incomplete (MUC5AC positive) IM glands (B and inset).

Figure 3- Expression of SOX2, CDX2, MUC5AC and MUC2 in gastric dysplasia. Dysplastic foci were immunostained for the gastric differentiation markers SOX2 (A and inset) and MUC5AC (D) and intestinal differentiation markers CDX2 (B) and MUC2 (D). In (A) arrowhead indicates dysplastic glands with SOX2 expression whereas the arrow indicates dysplastic glands without SOX2 expression. In (D) arrow indicates IM with homogeneous MUC2 expression as opposed to dysplasia where expression is scattered (arrowhead).

Figure 4- SOX2 overexpression in Caco-2 cells. Caco-2 colonic cell line was transiently transfected either with an empty vector or a SOX2 expression vector. The effect of SOX2 expression on CDX2 protein and RNA levels was assessed by western blotting (right) and qRT-PCR (left), respectively.

Figure 5- Schematic representation of gastric carcinogenesis.

Table 1 – Primary antibodies and immunohistochemistry conditions used in this study

Antibody	Clone	Antigen Retrieval Buffer	Antigen Retrieval Conditions	Dilution	Incubation time (min)	Localization	Source
CDX2	CDX2-88	Citrate Buffer 10mM pH6.0	40 minutes at 98°C	1:50	Overnight (4°C)	Nuclear	Biogenex, San Ramon, CA
MUC2	PMH1	0,1 U/mL Neuraminidase ^a	2h at 37°C	undiluted	Overnight (4°C)	Cytoplasmatic	Supernatant [27]
MUC5AC	CLH2	none	none	1:10	Overnight (4°C)	Cytoplasmatic	Supernatant [27]
MUC6	CLH5	none	none	1:10	Overnight (4°C)	Cytoplasmatic	Supernatant [27]
SOX2	SP-76	EDTA 10mM pH8.0	40 minutes at 98°C	1:50	1h (Room Temperature)	Nuclear	Cell Marque, Rockling, CA

^aNeuraminidase from *Clostridium perfringens* type VI (Sigma) was diluted in sodium acetate buffer (pH 5.5)

Table 2 - Expression of SOX2 in gastric IM.

Parameter	SOX2		<i>p</i> value
	Negative n (%)	Positive n (%)	
Complete IM foci (n=29)	27 (93%)	2 (7%)	<0.0001
Incomplete IM foci (n=26)	4 (15%)	22 (85%)	

Table 3 - Expression of SOX2, CDX2, MUC5AC, MUC2 and MUC6 in dysplasia

Parameter	Dysplasia (n=26)	
	Positive n (%)	Negative n (%)
SOX2	3 (12%)	23 (88%)
CDX2	20 (100%)	0 (0%)
MUC5AC	15 (58%)	11 (42%)
MUC2	12 (46%)	14 (54%)
MUC6^a	17 (68%)	8 (32%)

^aIn one case, MUC6 expression could not be evaluated

Figure 1




Figure 2




Figure 3




Figure 4





Figure 5



DISCUSSION

In spite of our well-established understanding of the phenotypic natural history occurring in the shift from native epithelia to invasive intestinal-type carcinoma in the gastric mucosa, the molecular typing of the precancerous changes in gastric mucosa remains elusive.¹ It is, however, becoming evident that the balance between gastric and intestinal differentiation may play a significant role, at least in the early steps of the carcinogenesis cascade. It has been shown that the transcription factors SOX2 and CDX2 are stomach and intestine-specific, respectively,^{2, 3} thus they may provide a useful tool to analyse this cascade.

In this study, we performed a systematic characterization of SOX2 expression along the precancerous cascade using immunohistochemistry in a series of 10 cases of normal gastric mucosa obtained from patients with no gastric pathology, 55 foci of IM adjacent to gastric carcinomas or to dysplasia, and 26 foci of dysplasia. We also used in vitro models to determine if there is a cross-regulation between SOX2 and CDX2.

Contrary to other authors, we have used an anti-SOX2 monoclonal antibody, raising the confidence level on the immunohistochemistry method and results. Also, the double immunostaining technique provides valuable information about the co-localization of the different markers. The small number of cases included might be pointed out as a limitation of the study.

We have shown that SOX2 expression is localized especially in the neck region of normal gastric glands, which is similar to other authors' findings.⁴ This is the proliferative compartment of the mucosa and the expected location of gastric stem cells,^{5, 6} which is consistent with previous papers describing SOX2 role in stem cell maintenance.⁷ However, the number of positive cells is greater than expected if SOX2 was only expressed in stem and progenitor cells, which suggests that this transcription

factor is not only required to switch on a gastric differentiation program, but also to maintain it in terminally differentiated cells.

In intestinal metaplasia (IM), SOX2 is expressed in cases that maintain a gastric phenotype, characterized by the presence of the MUC5AC mucin and defining the incomplete IM subtype. By contrast, in complete IM, SOX2 expression is lost concomitantly with loss of MUC5AC. These two aspects are especially evident with the double SOX2/MUC5AC immunostaining. Previous studies have also described SOX2 expression in human IM and, despite a different IM classification, the co-expression of SOX2 with MUC5AC or the absence of both SOX2 and MUC5AC was also observed.⁴

⁸ These observations are according with the expected role of SOX2 in the maintenance of gastric differentiation in incomplete type IM.

The expression pattern of SOX2 in gastric dysplasia was so far unknown. We showed for the first time that the majority of dysplastic lesions do not express SOX2 (3 out of 26) whereas all of them express CDX2. The positive cases were all low-grade dysplasias. One would be tempted to relate this finding with the previous reports of SOX2 possible tumour suppressor role,⁹ but the series of cases is short and it would be a risky shot. Despite our expectation of finding a relation between SOX2 and MUC5AC expression in dysplasia, we could not find such association nor with CDX2 expression in this lesion.

Although gastric carcinogenesis still is a mind-breaking issue, it is recognized that one of its major pathways develops in an *H. pylori*-induced background of chronic atrophic gastritis and is associated with reprogramming of differentiation leading to IM. CDX2 is documented as being a reliable responsible for this intestinal shift, leading to both types of IM. The differential expression of SOX2 in IM subtypes brought out the rationale for the maintenance of gastric differentiation in incomplete type IM.

While CDX2 expression is maintained in the progression to gastric dysplasia, SOX2 expression is substantially lost. These two events suggest that the progression from normal gastric mucosa to dysplasia occurs concomitantly with gain of intestinal and loss of gastric differentiation.

The regulatory mechanisms underlying CDX2 activation and SOX2 downregulation in the gastric setting are closer to be understood. A previous report of this group has shown that the BMP pathway, which is overexpressed in gastric IM, upregulates CDX2 and downregulates SOX2 in a gastric carcinoma cell line.^{10, 11} Furthermore, conditional knock-out of the CDX2 gene in mice intestines' showed increased SOX2 expression, with a consequent shift from an intestinal to a gastric/esophageal differentiation.^{12, 13} The same was observed when ectopic expression of SOX2 was induced in the intestine.¹⁴ These observations point to a mutual negative regulation of SOX2 and CDX2. In this study, we could not confirm the regulation of SOX2 by CDX2 by simply transfecting cell lines with CDX2, which suggests that co-factors or mediators are required for this process. We could though confirm SOX2 ability to negatively regulate CDX2 expression in one cell line.

One possible interpretation of our results would be that upon adverse events, the normal gastric mucosa evolves to a transient state of incomplete IM induced by CDX2. From this point, both SOX2 expression and gastric differentiation are lost, leading to complete IM with low evidence for further carcinogenic evolution. Otherwise, for unknown reasons, the transient incomplete IM expressing both transcription factors creates an unstable setting, prone to acquire genetic alterations in oncogenes or tumour suppressor genes that induces evolution to dysplasia, while maintaining the progressive loss of SOX2. This is also observed in gastric adenocarcinoma.^{9, 15}

In summary, SOX2 is expressed in the normal stomach and is maintained in incomplete IM explaining the maintenance of a gastric differentiation program in these

lesions. SOX2 expression is lost in the dysplastic lesions and in the complete IM subtype, so we hypothesize that incomplete IM may precede the onset of them both. The balance between gastric and intestinal differentiation has a significant impact in gastric carcinogenesis, at least in the early steps.

References

1. Fassan M, Baffa R, Kiss A. Advanced precancerous lesions within the GI tract: the molecular background. *Best Pract Res Clin Gastroenterol* 2013;**27**(2):159-69.
2. Almeida R, Silva E, Santos-Silva F, *et al.* Expression of intestine-specific transcription factors, CDX1 and CDX2, in intestinal metaplasia and gastric carcinomas. *J Pathol* 2003;**199**(1):36-40.
3. Tsukamoto T, Mizoshita T, Mihara M, *et al.* Sox2 expression in human stomach adenocarcinomas with gastric and gastric-and-intestinal-mixed phenotypes. *Histopathology* 2005;**46**(6):649-58.
4. Asonuma S, Imatani A, Asano N, *et al.* Helicobacter pylori induces gastric mucosal intestinal metaplasia through the inhibition of interleukin-4-mediated HMG box protein Sox2 expression. *Am J Physiol Gastrointest Liver Physiol* 2009;**297**(2):G312-22.
5. Pan Q, Nicholson AM, Barr H, *et al.* Identification of Lineage-Uncommitted, Long-Lived, Label-Retaining Cells in Healthy Human Esophagus and Stomach, and in Metaplastic Esophagus. *Gastroenterology* 2013;**144**(4):761-770.
6. Hoffmann W. Self-renewal of the gastric epithelium from stem and progenitor cells. *Front Biosci (Schol Ed)* 2013;**5**:720-31.
7. Sarkar A, Hochedlinger K. The Sox Family of Transcription Factors: Versatile Regulators of Stem and Progenitor Cell Fate. *Cell Stem Cell* 2013;**12**(1):15-30.
8. Tsukamoto T, Inada K, Tanaka H, *et al.* Down-regulation of a gastric transcription factor, Sox2, and ectopic expression of intestinal homeobox genes, Cdx1 and Cdx2: inverse correlation during progression from gastric/intestinal-mixed to complete intestinal metaplasia. *J Cancer Res Clin Oncol* 2004;**130**(3):135-45.
9. Otsubo T, Akiyama Y, Yanagihara K, *et al.* SOX2 is frequently downregulated in gastric cancers and inhibits cell growth through cell-cycle arrest and apoptosis. *Br J Cancer* 2008;**98**(4):824-31.

10. Camilo V, Barros R, Sousa S, *et al.* Helicobacter pylori and the BMP pathway regulate CDX2 and SOX2 expression in gastric cells. *Carcinogenesis* 2012;**33**(10):1985-92.
11. Barros R, Pereira B, Duluc I, *et al.* Key elements of the BMP/SMAD pathway co-localize with CDX2 in intestinal metaplasia and regulate CDX2 expression in human gastric cell lines. *J Pathol* 2008;**215**(4):411-20.
12. Stringer EJ, Duluc I, Saandi T, *et al.* Cdx2 determines the fate of postnatal intestinal endoderm. *Development* 2012;**139**(3):465-74.
13. Gao N, White P, Kaestner KH. Establishment of intestinal identity and epithelial-mesenchymal signaling by Cdx2. *Dev Cell* 2009;**16**(4):588-99.
14. Raghoebir L, Bakker ER, Mills JC, *et al.* SOX2 redirects the developmental fate of the intestinal epithelium toward a premature gastric phenotype. *J Mol Cell Biol* 2012;**4**(6):377-85.
15. Li XL, Eishi Y, Bai YQ, *et al.* Expression of the SRY-related HMG box protein SOX2 in human gastric carcinoma. *Int J Oncol* 2004;**24**(2):257-63.

ANEXO

**NORMAS DE PUBLICAÇÃO
“EUROPEAN JOURNAL OF CANCER”**

Introduction

The *European Journal of Cancer (EJC)* is an international multidisciplinary oncology journal, which publishes original research, reviews, and editorial comments on basic and preclinical cancer research, translational oncology, clinical oncology - including medical oncology, paediatric oncology, radiation oncology, and surgical oncology, and cancer epidemiology and prevention.

The *EJC* is the official journal of the [European Organisation for Research and Treatment of Cancer \(EORTC\)](#), the [European CanCER Organisation \(ECCO\)](#), the [European Association for Cancer Research \(EACR\)](#) and the [European Society of Breast Cancer Specialists \(EUSOMA\)](#).

Why publish your work in the EJC

Efficient peer-review: The *EJC*'s [Editors](#) and reviewers are committed to maintaining the high standards of the journal. All manuscripts submitted to the *EJC* will be evaluated by the journal's Editors. Some manuscripts may be rejected outright following this evaluation, allowing the authors to submit their work elsewhere. Those manuscripts which are judged as being eligible for consideration by the Editors will be subject to peer-review. The full peer-review policy for the *EJC* can be found [here](#).

The *EJC* has an [online submission](#) and peer-review system to facilitate the submission and peer-review of submitted manuscripts. Authors are able to upload their manuscript quickly and track the status of their manuscript throughout the review process.

Online Publication as an Article-in-Press: All manuscripts that are accepted for publication in the *EJC* are published online as Articles-in-Press within 4 - 5 weeks of acceptance. Articles-in-Press can be cited using the unique digital object identifier (DOI) which is assigned to all papers. In addition, Articles-in-Press are indexed in PubMed.

The *EJC*'s readership: The *EJC* is published in print and online and is available to more than 17,000 academic and clinical institutions, personal subscribers and members of the journal's Society Partners (the EORTC, ECCO, EACR and EUSOMA). In 2011 the journal had more than 83,000 articles downloaded per month from across the world.

The *EJC* has an iPad app, making each issue of the journal available to our readers in this format. Visit <http://itunes.apple.com/us/app/ejc-app/id461022524?mt=8> to find out more.

eTables of Contents: Our readers can elect to receive free eTables of Contents via email, which are sent out for each issue of the journal.

Abstracting and Indexing Coverage: The *EJC* is included in the leading abstracting and indexing services, including PubMed/MEDLINE, Current Contents/Clinical Medicine, the Science Citation Index, the Journal Citation Reports and Scopus.

HINARI: The *EJC* is part of HINARI (Health InterNetwork Access to Research Initiative) administered by the [World Health Organisation](#). The HINARI Programme enables developing countries to gain access to one of the world's largest collections of biomedical and health literature.

Why read the EJC

High quality original research and review articles: The *EJC* is a multidisciplinary journal, publishing high quality original research and review articles on basic and preclinical research, translational oncology, clinical oncology (medical, paediatric, radiation, surgical), and cancer epidemiology and prevention from across the world.

Guidelines: The *EJC* is privileged to publish selected treatment guidelines from the journal's Society Partners (the EORTC, ECCO, EACR and EUSOMA) and other groups.

Special Series: The *EJC* publishes special series of papers on linked topics of particular relevance and interest to the journal's readers.

***EJC* News and *EJC* News Focus:** Each month, the journal publishes the *EJC* News, discussing and commenting on current issues in cancer, together with the *EJC* News Focus, a series of video interviews with key opinion leaders in oncology.

Types of paper

The *EJC* will consider the following types of manuscript for publication:

Editorial comments: Editorial comments are generally invited by the *EJC*'s Editorial Team. They are 1,500 words in length with no abstract or keywords.

Original research articles: Original research articles have a limit of 2,500 words and no more than 40 references. Authors are asked to provide a structured abstract of no more than 250 words and a list of keywords.

Review articles: Review articles have a limit of 3,000 words with an unlimited number of references. Authors are asked to provide an unstructured abstract of no more than 250 words and a list of keywords.

Current Perspectives: Current Perspectives have a limit of 2,000 words with an unlimited number of tables and figures and a limit of 30 references. Authors should provide an unstructured abstract of no more than 250 words and a list of keywords.

Letters to the Editor: Letters to the Editor which comment directly on work that has previously been published in the EJC will be considered. Letters have a limit of 1,000 words and should be structured as a Letter, without an abstract or keywords. The author of the paper on which the Letter is commenting will be contacted and invited to respond.

Contact details for submission

Manuscripts and the accompanying [Author Form](#) should be submitted online at <http://ees.elsevier.com/ejc> and the instructions on the site should be closely followed. Authors may submit manuscripts and track their progress to final decision. Reviewers can download manuscripts and submit their reports to the Editors.

The full contact details for the Editorial Office are shown below:

European Journal of Cancer
Elsevier Ltd
The Boulevard, Langford Lane
Oxford OX5 1GB
UK
Tel : +44 (0)1865 843282
Fax : +44 (0)1865 843977
Email : ejcancer@elsevier.com

In a covering letter, please identify the person responsible for editorial correspondence (address, telephone and fax numbers and e-mail address).

Before you begin

Ethics in publishing

For information on Ethics in publishing and Ethical guidelines for journal publication see ➡ <http://www.elsevier.com/publishingethics> and ➡ <http://www.elsevier.com/journal-authors/ethics>.

Human and animal rights

If the work involves the use of animal or human subjects, the author should ensure that the work described has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans ➡ <http://www.wma.net/en/30publications/10policies/b3/index.html>; EU Directive 2010/63/EU for animal experiments ➡ http://ec.europa.eu/environment/chemicals/lab_animals/legislation_en.htm; Uniform Requirements for manuscripts submitted to Biomedical journals ➡ <http://www.icmje.org>. Authors should include a statement in the manuscript that informed consent was obtained for experimentation with human subjects. The privacy rights of human subjects must always be observed.

Conflict of interest

All authors must disclose any financial and personal relationships with other people or organizations that could inappropriately influence (bias) their work. Examples of potential conflicts of interest include employment, consultancies, stock ownership, honoraria, paid expert testimony, patent applications/registrations, and grants or other funding. See also ➡ <http://www.elsevier.com/conflictsofinterest>. Further information and an example of a Conflict of Interest form can be found at: ➡ http://help.elsevier.com/app/answers/detail/a_id/286/p/7923.

Submission declaration

Submission of an article implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see ➡ <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere including electronically in the same form, in English or in any other language, without the written consent of the copyright-holder.

Author form

The corresponding author must submit a completed [Author Form](#) with their submission. The form enables the corresponding author to record the contribution of each author and confirms whether ethical approval for the study has been sought, whether funding for the study was provided and whether there are any conflicts of interest. It must be signed by the corresponding author and uploaded to EES with the manuscript.

Reporting clinical trials

Randomized controlled trials should be presented according to the CONSORT guidelines. At manuscript submission, authors must provide the CONSORT checklist accompanied by a flow diagram that illustrates the progress of patients through the trial, including recruitment, enrollment, randomization, withdrawal and completion, and a detailed description of the randomization procedure. The CONSORT checklist and template flow diagram can be found on ➡ <http://www.consort-statement.org>.

Registration of clinical trials

Registration in a public trials registry is a condition for publication of clinical trials in this journal in accordance with International Committee of Medical Journal Editors (ICMJE, <http://www.icmje.org>) recommendations. Trials must register at or before the onset of patient enrolment. The clinical trial registration number should be included at the end of the abstract of the article. A clinical trial is defined as any research study that prospectively assigns human participants or groups of humans to one or more health-related interventions to evaluate the effects of health outcomes. Health-related interventions include any intervention used to modify a biomedical or health-related outcome (for example drugs, surgical procedures, devices, behavioural treatments, dietary interventions, and process-of-care changes). Health outcomes include any biomedical or health-related measures obtained in patients or participants, including pharmacokinetic measures and adverse events. Purely observational studies (those in which the assignment of the medical intervention is not at the discretion of the investigator) will not require registration.

Copyright

This journal offers authors a choice in publishing their research: Open Access and Subscription.

For Subscription articles

Upon acceptance of an article, authors will be asked to complete a 'Journal Publishing Agreement' (for more information on this and copyright, see <http://www.elsevier.com/copyright>). An e-mail will be sent to the corresponding author confirming receipt of the manuscript together with a 'Journal Publishing Agreement' form or a link to the online version of this agreement. Subscribers may reproduce tables of contents or prepare lists of articles including abstracts for internal circulation within their institutions. Permission of the Publisher is required for resale or distribution outside the institution and for all other derivative works, including compilations and translations (please consult <http://www.elsevier.com/permissions>). If excerpts from other copyrighted works are included, the author(s) must obtain written permission from the copyright owners and credit the source(s) in the article. Elsevier has preprinted forms for use by authors in these cases: please consult <http://www.elsevier.com/permissions>.

For Open Access articles

Upon acceptance of an article, authors will be asked to complete an 'Exclusive License Agreement' (for more information see <http://www.elsevier.com/OAauthoragreement>). Permitted reuse of open access articles is determined by the author's choice of user license (see <http://www.elsevier.com/openaccesslicenses>).

Retained author rights

As an author you (or your employer or institution) retain certain rights. For more information on author rights for: Subscription articles please see <http://www.elsevier.com/journal-authors/author-rights-and-responsibilities>. Open access articles please see <http://www.elsevier.com/OAauthoragreement>.

Role of the funding source

You are requested to identify who provided financial support for the conduct of the research and/or preparation of the article and to briefly describe the role of the sponsor(s), if any, in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. If the funding source(s) had no such involvement then this should be stated. Please see <http://www.elsevier.com/funding>.

Microarray data

The data should be made available to our readers/reviewers through a MIAME compliant online database or at least a website address.

Biomarkers

Submissions which report biomarkers must comply with the REMARK Guidelines (McShane L, Altman DG, Sauerbrei W, Taube SE, Gion M, Clark GM for the Statistics Subcommittee of the NCI-EORTC Working Group on Cancer diagnostics. Reporting recommendations for tumor MARKer prognostic studies (REMARK), *Eur J Cancer*, 2005, 41, 1690-1696.)

Funding body agreements and policies

Elsevier has established agreements and developed policies to allow authors whose articles appear in journals published by Elsevier, to comply with potential manuscript archiving requirements as specified as conditions of their grant awards. To learn more about existing agreements and policies please visit <http://www.elsevier.com/fundingbodies>.

Open access

This journal offers authors a choice in publishing their research:

Open Access

- Articles are freely available to both subscribers and the wider public with permitted reuse
- An Open Access publication fee is payable by authors or their research funder

Subscription

- Articles are made available to subscribers as well as developing countries and patient groups through our access programs (



- No Open Access publication fee


All articles published Open Access will be immediately and permanently free for everyone to read and download. Permitted reuse is defined by your choice of one of the following Creative Commons user licenses: **Creative Commons Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)**: for non-commercial purposes, lets others distribute and copy the article, to create extracts, abstracts and other revised versions, adaptations or derivative works of or from an article (such as a translation), to include in a collective work (such as an anthology), to text and data mine the article, as long as they credit the author(s), do not represent the author as endorsing their adaptation of the article, do not modify the article in such a way as to damage the author's honor or reputation, and license their new adaptations or creations under identical terms (CC BY-NC-SA).

Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND): for non-commercial purposes, lets others distribute and copy the article, and to include in a collective work (such as an anthology), as long as they credit the author(s) and provided they do not alter or modify the article.



Elsevier has established agreements with funding bodies,  <http://www.elsevier.com/fundingbodies>. This ensures authors can comply with funding body Open Access requirements, including specific user licenses, such as CC BY. Some authors may also be reimbursed for associated publication fees. If you need to comply with your funding body policy, you can apply for the CC BY license after your manuscript is accepted for publication.

To provide Open Access, this journal has a publication fee which needs to be met by the authors or their research funders for each article published Open Access.

Your publication choice will have no effect on the peer review process or acceptance of submitted articles.

The publication fee for this journal is **\$3,000**, excluding taxes. Learn more about Elsevier's pricing policy:  <http://www.elsevier.com/openaccesspricing>.

Language (usage and editing services)

Please write your text in good English (American or British usage is accepted, but not a mixture of these). Authors who feel their English language manuscript may require editing to eliminate possible grammatical or spelling errors and to conform to correct scientific English may wish to use the English Language Editing service available from Elsevier's WebShop ( <http://webshop.elsevier.com/languageediting/>) or visit our customer support site ( <http://support.elsevier.com>) for more information.

Submission

Submission to this journal proceeds totally online and you will be guided stepwise through the creation and uploading of your files. The system automatically converts source files to a single PDF file of the article, which is used in the peer-review process. Please note that even though manuscript source files are converted to PDF files at submission for the review process, these source files are needed for further processing after acceptance. All correspondence, including notification of the Editor's decision and requests for revision, takes place by e-mail removing the need for a paper trail.

Submit your article

Please submit your article via  <http://www.ees.elsevier.com/ejc>.

Referees

Please submit, with the manuscript, the names, addresses and e-mail addresses of five potential referees. Note that the editor retains the sole right to decide whether or not the suggested reviewers are used.

Preparation

Article structure

Please write your text in good English with double line-spacing. Original research manuscripts should be no more than 2,500 words in length, with a maximum of 40 references, and review articles should be no more than 3,000 words in length, with an unlimited number of references. Abstracts should not exceed 250 words in length. Manuscripts may be altered to meet the EJC's style.

Manuscripts containing research data generally follow the order: Introduction, Patients (or Materials) and Methods, Results, and Discussion.

Appendices

If there is more than one appendix, they should be identified as A, B, etc. Formulae and equations in appendices should be given separate numbering: Eq. (A.1), Eq. (A.2), etc.; in a subsequent appendix, Eq. (B.1) and so on. Similarly for tables and figures: Table A.1; Fig. A.1, etc.

Essential title page information

- **Title.** Concise and informative. Titles are often used in information-retrieval systems. Avoid abbreviations and formulae where possible.
- **Author names and affiliations.** Where the family name may be ambiguous (e.g., a double name), please indicate this clearly. Present the authors' affiliation addresses (where the actual work was done) below the names. Indicate all affiliations with a lower-case superscript letter immediately after the author's name and in front of the appropriate address. Provide the full postal address of each affiliation, including the country name and, if available, the e-mail address of each author.
- **Corresponding author.** Clearly indicate who will handle correspondence at all stages of refereeing and publication, also post-publication. **Ensure that phone numbers (with country and area code) are provided in addition to the e-mail address and the complete postal address. Contact details must be kept up to date by the corresponding author.**
- **Present/permanent address.** If an author has moved since the work described in the article was done, or was visiting at the time, a 'Present address' (or 'Permanent address') may be indicated as a footnote to that author's name. The address at which the author actually did the work must be retained as the main, affiliation address. Superscript Arabic numerals are used for such footnotes.

The title page should also include details of sources of support in the form of grants, equipment, and drugs.

Abstract

The second page should start with the abstract, which should be up to a maximum of 250 words and must include the aim of the study, a brief summary of the methods, results and a concluding statement.

Graphical abstract

A Graphical abstract is optional and should summarize the contents of the article in a concise, pictorial form designed to capture the attention of a wide readership online. Authors must provide images that clearly represent the work described in the article. Graphical abstracts should be submitted as a separate file in the online submission system. Image size: Please provide an image with a minimum of 531 × 1328 pixels (h × w) or proportionally more. The image should be readable at a size of 5 × 13 cm using a regular screen resolution of 96 dpi. Preferred file types: TIFF, EPS, PDF or MS Office files. See <http://www.elsevier.com/graphicalabstracts> for examples.

Authors can make use of Elsevier's Illustration and Enhancement service to ensure the best presentation of their images also in accordance with all technical requirements: [Illustration Service](#).

Highlights

Highlights are mandatory for this journal. They consist of a short collection of bullet points that convey the core findings of the article and should be submitted in a separate file in the online submission system. Please use 'Highlights' in the file name and include 3 to 5 bullet points (maximum 85 characters, including spaces, per bullet point). See <http://www.elsevier.com/highlights> for examples.

Keywords

Include up to 10 key words from the Medical Subject Headings from *Index Medicus*.

Acknowledgements

All contributors who do not meet the criteria for authorship as defined by the ICMJE should be listed in an acknowledgments section. Examples of those who might be acknowledged include a person who provided purely technical help, writing assistance, or a department chair who provided only general support. Authors should disclose whether they had any writing assistance and identify the entity that paid for this assistance.

Footnotes

Footnotes should be used sparingly. Number them consecutively throughout the article, using superscript Arabic numbers. Many wordprocessors build footnotes into the text, and this feature may be used. Should this not be the case, indicate the position of footnotes in the text and present the footnotes themselves separately at the end of the article. Do not include footnotes in the Reference list.

Table footnotes

Indicate each footnote in a table with a superscript lowercase letter.

Artwork

Electronic artwork

General points

- Make sure you use uniform lettering and sizing of your original artwork.
- Embed the used fonts if the application provides that option.
- Aim to use the following fonts in your illustrations: Arial, Courier, Times New Roman, Symbol, or use fonts that look similar.
- Number the illustrations according to their sequence in the text.
- Use a logical naming convention for your artwork files.
- Provide captions to illustrations separately.
- Size the illustrations close to the desired dimensions of the printed version.

- Submit each illustration as a separate file.

A detailed guide on electronic artwork is available on our website:



<http://www.elsevier.com/artworkinstructions>

You are urged to visit this site; some excerpts from the detailed information are given here.

Formats

If your electronic artwork is created in a Microsoft Office application (Word, PowerPoint, Excel) then please supply 'as is' in the native document format.

Regardless of the application used other than Microsoft Office, when your electronic artwork is finalized, please 'Save as' or convert the images to one of the following formats (note the resolution requirements for line drawings, halftones, and line/halftone combinations given below):

EPS (or PDF): Vector drawings, embed all used fonts.

TIFF (or JPEG): Color or grayscale photographs (halftones), keep to a minimum of 300 dpi.


TIFF (or JPEG): Bitmapped (pure black & white pixels) line drawings, keep to a minimum of 1000 dpi.

TIFF (or JPEG): Combinations bitmapped line/half-tone (color or grayscale), keep to a minimum of 500 dpi.

Please do not:

- Supply files that are optimized for screen use (e.g., GIF, BMP, PICT, WPG); these typically have a low number of pixels and limited set of colors;
- Supply files that are too low in resolution;
- Submit graphics that are disproportionately large for the content.

Color artwork

Please make sure that artwork files are in an acceptable format (TIFF (or JPEG), EPS (or PDF), or MS Office files) and with the correct resolution. If, together with your accepted article, you submit usable color figures then Elsevier will ensure, at no additional charge, that these figures will appear in color on the Web (e.g., ScienceDirect and other sites) regardless of whether or not these illustrations are reproduced in color in the printed version. **For color reproduction in print, you will receive information regarding the costs from Elsevier after receipt of your accepted article.** Please indicate your preference for color: in print or on the Web only. For further information on the preparation of electronic artwork, please see  <http://www.elsevier.com/artworkinstructions>.

Please note: Because of technical complications which can arise by converting color figures to 'gray scale' (for the printed version should you not opt for color in print) please submit in addition usable black and white versions of all the color illustrations.

Illustration services


Elsevier's WebShop ( <http://webshop.elsevier.com/illustrationservices>) offers Illustration Services to authors preparing to submit a manuscript but concerned about the quality of the images accompanying their article. Elsevier's expert illustrators can produce scientific, technical and medical-style images, as well as a full range of charts, tables and graphs. Image 'polishing' is also available, where our illustrators take your image(s) and improve them to a professional standard. Please visit the website to find out more.

Figure captions

Ensure that each illustration has a caption. Supply captions separately, not attached to the figure. A caption should comprise a brief title (**not** on the figure itself) and a description of the illustration. Keep text in the illustrations themselves to a minimum but explain all symbols and abbreviations used.

Tables

Number tables consecutively in accordance with their appearance in the text. Place footnotes to tables below the table body and indicate them with superscript lowercase letters. Avoid vertical rules. Be sparing in the use of tables and ensure that the data presented in tables do not duplicate results described elsewhere in the article.

References

Citation in text

Please ensure that every reference cited in the text is also present in the reference list (and vice versa). Any references cited in the abstract must be given in full. Unpublished results and personal communications are not recommended in the reference list, but may be mentioned in the text. If these references are included in the reference list they should follow the standard reference style of the journal and should include a substitution of the publication date with either 'Unpublished results' or 'Personal communication'. Citation of a reference as 'in press' implies that the item has been accepted for publication.

Reference formatting

There are no strict requirements on reference formatting at submission. References can be in any style or format as long as the style is consistent. Where applicable, author(s) name(s), journal title/book title, chapter title/article title, year of publication, volume number/book chapter and the pagination must be present. Use of DOI is highly encouraged. The reference style used

by the journal will be applied to the accepted article by Elsevier at the proof stage. Note that missing data will be highlighted at proof stage for the author to correct. If you do wish to format the references yourself they should be arranged according to the following examples:

Revised manuscripts

Authors who have been asked to revise their manuscript by the Editors should submit a file which clearly shows the changes that have been made via the 'track changes' function or text highlighting, and a clean copy of the revised manuscript (i.e. a file which contains the changes without any highlighting). Authors are requested **not** to upload a copy of the original manuscript.

Reference style

Text: Indicate references by number(s) in square brackets in line with the text. The actual authors can be referred to, but the reference number(s) must always be given.

List: Number the references (numbers in square brackets) in the list in the order in which they appear in the text.

Examples:

Reference to a journal publication:

[1] Van der Geer J, Hanraads JAJ, Lupton RA. The art of writing a scientific article. *J Sci Commun* 2010;163:51–9.

Reference to a book:

[2] Strunk Jr W, White EB. *The elements of style*. 4th ed. New York: Longman; 2000.

Reference to a chapter in an edited book:

[3] Mettam GR, Adams LB. How to prepare an electronic version of your article. In: Jones BS, Smith RZ, editors. *Introduction to the electronic age*, New York: E-Publishing Inc; 2009, p. 281–304.

Note shortened form for last page number. e.g., 51–9, and that for more than 6 authors the first 6 should be listed followed by 'et al.' For further details you are referred to 'Uniform Requirements for Manuscripts submitted to Biomedical Journals' (*J Am Med Assoc* 1997;277:927–34) (see also http://www.nlm.nih.gov/bsd/uniform_requirements.html).

AudioSlides

The journal encourages authors to create an AudioSlides presentation with their published article. AudioSlides are brief, webinar-style presentations that are shown next to the online article on ScienceDirect. This gives authors the opportunity to summarize their research in their own words and to help readers understand what the paper is about. More information and examples are available at <http://www.elsevier.com/audioslides>. Authors of this journal will automatically receive an invitation e-mail to create an AudioSlides presentation after acceptance of their paper.

Supplementary data

Elsevier accepts electronic supplementary material to support and enhance your scientific research. Supplementary files offer the author additional possibilities to publish supporting applications, high-resolution images, background datasets, sound clips and more. Supplementary files supplied will be published online alongside the electronic version of your article in Elsevier Web products, including ScienceDirect: <http://www.sciencedirect.com>. In order to ensure that your submitted material is directly usable, please provide the data in one of our recommended file formats. Authors should submit the material in electronic format together with the article and supply a concise and descriptive caption for each file. For more detailed instructions please visit our artwork instruction pages at <http://www.elsevier.com/artworkinstructions>.

Submission checklist

The following list will be useful during the final checking of an article prior to sending it to the journal for review. Please consult this Guide for Authors for further details of any item.

Ensure that the following items are present:

One author has been designated as the corresponding author with contact details:

- E-mail address
- Full postal address
- Phone numbers

All necessary files have been uploaded, and contain:

- Keywords
- All figure captions
- All tables (including title, description, footnotes)

Further considerations

- Manuscript has been 'spell-checked' and 'grammar-checked'
 - References are in the correct format for this journal
 - All references mentioned in the Reference list are cited in the text, and vice versa
 - Permission has been obtained for use of copyrighted material from other sources (including the Web)
 - Color figures are clearly marked as being intended for color reproduction on the Web (free of charge) and in print, or to be reproduced in color on the Web (free of charge) and in black-and-white in print
 - If only color on the Web is required, black-and-white versions of the figures are also supplied for printing purposes
- For any further information please visit our customer support site at <http://support.elsevier.com>.

After Acceptance

Use of the Digital Object Identifier

The Digital Object Identifier (DOI) may be used to cite and link to electronic documents. The DOI consists of a unique alphanumeric character string which is assigned to a document by the publisher upon the initial electronic publication. The assigned DOI never changes. Therefore, it is an ideal medium for citing a document, particularly 'Articles in press' because they have not yet received their full bibliographic information. Example of a correctly given DOI (in URL format; here an article in the journal *Physics Letters B*):



<http://dx.doi.org/10.1016/j.physletb.2010.09.059>



When you use a DOI to create links to documents on the web, the DOIs are guaranteed never to change.

Online proof correction





Corresponding authors will receive an e-mail with a link to our online proofing system, allowing annotation and correction of proofs online. The environment is similar to MS Word: in addition to editing text, you can also comment on figures/tables and answer questions from the Copy Editor. Web-based proofing provides a faster and less error-prone process by allowing you to directly type your corrections, eliminating the potential introduction of errors.

If preferred, you can still choose to annotate and upload your edits on the PDF version. All instructions for proofing will be given in the e-mail we send to authors, including alternative methods to the online version and PDF. We will do everything possible to get your article published quickly and accurately - please upload all of your corrections within 48 hours. It is important to ensure that all corrections are sent back to us in one communication. Please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility. Note that Elsevier may proceed with the publication of your article if no response is received.

Offprints

The corresponding author, at no cost, will be provided with a PDF file of the article via e-mail (the PDF file is a watermarked version of the published article and includes a cover sheet with the journal cover image and a disclaimer outlining the terms and conditions of use). For an extra charge, paper offprints can be ordered via the offprint order form which is sent once the article is accepted for publication. Both corresponding and co-authors may order offprints at any time via Elsevier's WebShop ( <http://webshop.elsevier.com/myarticleservices/offprints>). Authors requiring printed copies of multiple articles may use Elsevier WebShop's 'Create Your Own Book' service to collate multiple articles within a single cover ( <http://webshop.elsevier.com/myarticleservices/offprints/myarticleservices/booklets>).

Author Inquiries

For inquiries relating to the submission of articles (including electronic submission) please visit this journal's homepage. For detailed instructions on the preparation of electronic artwork, please visit  <http://www.elsevier.com/artworkinstructions>. Contact details for questions arising after acceptance of an article, especially those relating to proofs, will be provided by the publisher. You can track accepted articles at  <http://www.elsevier.com/trackarticle>. You can also check our Author FAQs at  <http://www.elsevier.com/authorFAQ> and/or contact Customer Support via  <http://support.elsevier.com>.